

Protocol version 2020_3_14

COVID treatment protocol



EXCEPTIONAL CARE. WITHOUT EXCEPTION.
© 2020 Boston Medical Center

- No steroids for mild/moderate sickness
- Avoid Nebs
- Steroid use should be a discussion between ID and Pulmonary/Critical Care

All Persons Under Investigation

Contact/Droplet Precautions
 Comprehensive Respiratory Panel *
 COVID-19 PCR*
 Sputum Cultures*
 MRSA Nares*
 Procalcitonin
 G6PD
 Imaging: CXR, EKG
 Continuous Pulse OX

*NOTE: N95 respirator specimen collection. Collect all at same time to minimize exposure

Hydroxychloroquine dosing:
 400mg BID first day then
 200mg BID for 4 days (total 5d)

Remdesivir dosing:
 200mg first day then
 100mg daily for 4 days (total 5d)

COVID PCR Negative
AND
Alternative Diagnosis

Stop Hydroxychloroquine

Temp > 101F
AND O2 sat <90%
 Start Hydroxychloroquine
 Trend QTc qD
 Trend Glucose qD

COVID PCR Positive

Call Infectious Diseases
 Continue Hydroxychloroquine
 Request Remdesivir through
 online portal:

<https://rdvcu.gilead.com>